

**Physiotherapy works ✓**

# Vestibular rehabilitation

**Vestibular rehabilitation (VR) is effective in reducing dizziness and vertigo and associated falls and improving quality of life in people with disorders involving balance mechanisms**

Good balance is essential for daily life, from getting out of bed to crossing the road. A healthy balance system uses information from the brain, inner ears, eyes, and joints, and enables people to see clearly when moving their head.<sup>(1)</sup>

Balance disorders and dizziness are a growing public health concern across all age groups. Prevalence increases with age and these disorders are some of the most common complaints reported to GPs by older patients.<sup>(1)</sup> Untreated, they result in worsening unsteadiness, loss of confidence and anxiety, difficulty with concentration and memory, inactivity, loss of independence, an increased risk of falls and costly falls-related injuries.<sup>(1)(2)</sup>

## What causes dizziness?

The most common form of dizziness and balance disorder is BPPV (Benign Paroxysmal Positional Vertigo), a disorder of the inner ear resulting in episodes of intense dizziness on certain head movements.

Other causes include side effects from medication or alcohol, brain disorders such as stroke, concussion or cerebellar degeneration, cardiac problems and anxiety.

**VR -  
a successful  
treatment  
for people with  
common balance  
and dizziness  
disorders**

## What is vestibular rehabilitation

VR is an exercise-based programme to encourage the central nervous system to compensate for problems in the inner ear. Following thorough examination, specialist physiotherapists develop personalised exercise-based management plans including home exercise programmes. ▶▶

## Size of the problem



- Almost 1 in 4 adults under 65 years report dizziness or vertigo, often causing occupational difficulties or preventing employment but less than **25 per cent** had received treatment<sup>(10)</sup>
- About **20 per cent** with vestibular vertigo avoid leaving the house and **41 per cent** require sick leave<sup>(7)</sup>
- Over **38 per cent** of those aged over 85 years fall annually; half in combination with dizziness or blackouts
- **2.4 per cent** lifetime prevalence of BPPV alone, with only **8 per cent** receiving effective treatment.<sup>(8-10)</sup>





## Cost of ill health

- **80 per cent** of unexplained fallers attending A&E had possible vestibular loss, **41 per cent** had vertigo, with **30-40 per cent** admitted to hospital<sup>(11)</sup>
- Fallers use more than 4 million bed-days, cost the NHS **£2.3 billion** and incur total costs including social-care of **£5.6 billion**.<sup>(1,2)</sup> Costs per faller are **£1,720**, rising to **£8,600** for those seeking medical care<sup>(1,3)</sup>
- The costs of BPPV are significant; American healthcare costs to diagnosis BPPV alone approach **\$2 billion**<sup>(14)</sup>
- Untreated dizziness/balance disorders may lead to anxiety, avoidance strategies and increased risk of falling, impacting further on resources.<sup>(8, 15)</sup>

### A management plan may include:

- Eye and head movement exercises<sup>(2)</sup>
- Exercises to improve standing balance and walking<sup>(3)</sup>
- Specific techniques to reposition fragments in the inner ear which cause dizziness in BPPV<sup>(2)</sup>
- Education and advice on activities of daily living to improve confidence & function
- Advice on falls reduction/prevention.

### Significant improvement is achieved in:

- 80 per cent of patients with unilateral vestibular loss (e.g. labyrinthitis, acoustic neuroma surgery)<sup>(3)</sup>
- 80 per cent of patients with BPPV after one treatment session<sup>(4)</sup>
- Patients with a central cause (e.g. stroke) may show some improvement
- Use of VR in concussion,<sup>(5)</sup> migraine,<sup>(6)</sup> and non-specific balance loss of the elderly may be helpful.

Physiotherapists are ideally placed to provide early screening and intervention to minimise the impact on daily life of balance and dizziness disorders.

Specialist physiotherapists provide personalised VR programmes which are clinically effective reducing distressing symptoms and improving quality of life.

### Case study

Over a six month period, the introduction of an award-winning Specialist Physiotherapy and Audiology service for patients referred with balance/dizziness disorders at Aintree Hospitals, Liverpool, has reduced waiting times and achieved excellent patient satisfaction. 80 per cent of patients are now managed exclusively in physiotherapy led clinics with 52 per cent assessed, diagnosed and treated in one session.



### Further information

#### CSP Enquiry Handling Unit

**Tel:** 0207 306 6666  
**Email:** enquiries@csp.org.uk  
**Web:** www.csp.org.uk



### Acknowledgements

Nova Mullin, Specialist Physiotherapist Balance and Dizziness  
 @Aintree University Hospital NHS Foundation Trust, Liverpool.

## References

1. Sturme DL, St George R, Lord SR. Balance disorders in the elderly. *Neurophysiol Clin.* 2008 Dec;38(6):467-78.
2. Bhattacharyya N, Baugh RF, Orvidas L, et al. Clinical practice guideline: benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg.* 2008 Nov;139(5 Suppl 4):S47-81. URL: [http://oto.sagepub.com/content/139/5\\_suppl/S47.full.pdf](http://oto.sagepub.com/content/139/5_suppl/S47.full.pdf)
3. Hillier Susan L, McDonnell M. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database of Systematic Reviews.* 2011(2):Art. No.: CD005397. <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD005397.pub3/abstract>
4. Epley JM. The canalith repositioning procedure: for treatment of benign paroxysmal positional vertigo. *Otolaryngol Head Neck Surg.* 1992 Sep;107(3):399-404.
5. Alsalaheen BA, Mucha A, Morris LO, et al. Vestibular rehabilitation for dizziness and balance disorders after concussion. *Journal of Neurologic Physical Therapy: JNPT.* 2010 Jun;34(2):87-93.
6. Gottshall KR, Moore RJ, Hoffer ME. Vestibular rehabilitation for migraine-associated dizziness. *The International Tinnitus Journal.* 2005;11(1):81-4.
7. Neuhauser HK, Radtke A, von Brevern M, et al. Burden of dizziness and vertigo in the community. *Archives of Internal Medicine.* 2008;168(19):2118-24. URL: <http://dx.doi.org/10.1001/archinte.168.19.2118>
8. von Brevern M, Radtke A, Lezius F, et al. Epidemiology of benign paroxysmal positional vertigo: a population based study. *Journal of Neurology, Neurosurgery, and Psychiatry.* 2007 Jul;78(7):710-5. URL: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2117684/pdf/710.pdf>
9. Stevens KN, Lang IA, Guralnik JM, et al. Epidemiology of balance and dizziness in a national population: findings from the English Longitudinal Study of Ageing. *Age and Ageing.* 2008 May 1, 2008;37(3):300-5. URL: <http://ageing.oxfordjournals.org/content/37/3/300.abstract>
10. Collerton J, Kingston A, Bond J, et al. The personal and health service impact of falls in 85 year olds: cross-sectional findings from the Newcastle 85+ cohort study. *PLoS one.* 2012;7(3):e33078. URL: [www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0033078](http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0033078)
11. Pothula VB, Chew F, Lesser TH, et al. Falls and vestibular impairment. *Clinical Otolaryngology and Allied Sciences.* 2004 Apr;29(2):179-82. URL: <http://onlinelibrary.wiley.com/doi/10.1111/j.0307-7772.2004.00785.x/abstract>
12. Tian Y, Thompson J, Buck D, et al. Exploring the system-wide costs of falls in older people in Torbay. London: The King's Fund; 2013. URL: <http://www.kingsfund.org.uk/publications/exploring-system-wide-costs-falls-older-people-torbay>
13. Craig J, Murray A, Mitchell S, et al. The high cost to health and social care of managing falls in older adults living in the community in Scotland. *Scottish Medical Journal.* [in press]
14. Neuhauser HK. Epidemiology of vertigo. *Current Opinion in Neurology.* 2007 Feb;20(1):40-6.
15. Yardley L, Redfern MS. Psychological factors influencing recovery from balance disorders. *Journal of Anxiety Disorders.* 2001;15(1):107-19.